

the interference between atoms in those two momentum states that produces density modulations.

To observe the onset of the ordered phase, they gradually increased the pump power while monitoring the light leaking out of the cavity. As long as the power remained below a critical threshold, no photons were detected outside the cavity. But above that threshold they noticed an abrupt increase in the number of photons. The time-of-flight images shown in figure 2 capture the change in symmetry in the momentum distribution of the BEC.

A fortunate aspect of the experiment is that the cavity output reveals time-resolved information. One experimental challenge is to probe the system's

dynamics close to the phase transition.

A new paradigm

The study of a BEC in an optical cavity represents a departure from most condensed-matter experiments, which typically probe short-range interactions among atoms. Because every atom feels the presence of every other atom via the mediating superradiant light field, the atomic interactions are truly long range: The number of nearest neighbors is effectively equal to the number of particles.

Many-body cavity-QED experiments may also provide a new perspective on phase transitions, conventionally studied in closed systems at equilibrium. Esslinger's BEC remains in a stable ground state throughout the ex-

periment. But the system is open, externally driven, and dissipative—far from what Dicke originally considered. That his Hamiltonian quantitatively captures the essential physics is thus all the more remarkable.

Mark Wilson

References

1. R. H. Dicke, *Phys. Rev.* **93**, 99 (1954).
2. S. Inouye, *Science* **285**, 571 (1999).
3. K. Baumann et al., *Nature* **464**, 1301 (2010).
4. A. T. Black, H. W. Chan, V. Vuletić, *Phys. Rev. Lett.* **91**, 203001 (2003).
5. P. Domokos, H. Ritsch, *Phys. Rev. Lett.* **89**, 253003 (2002).
6. F. Brennecke et al., *Nature* **450**, 268 (2007); Y. Colombe et al., *Nature* **450**, 272 (2007).
7. F. Dimer et al., *Phys. Rev. A* **75**, 013804 (2007).

A complex fluid exhibits unexpected heterogeneous flow

Depending on the conditions, the localized flow can persist for minutes, hours, or more than a day.

Flowing water, like other Newtonian fluids, is fully described by the Navier-Stokes equations. But many everyday materials, such as foams, emulsions, and colloids, are complex fluids that lack a description of similar generality. Faced with that gap, researchers instead look for similarities among different fluid systems to better classify their behavior. One such class, the yield stress fluids (YSFs), includes materials such as mayonnaise, hair gel, and toothpaste that hold their shape under low stress but flow under high stress.

Recently, YSFs were further classified into those that exhibit thixotropy—the decrease of viscosity with time during continued flow—and those that don't.¹ Thixotropy in a YSF leads to heterogeneous flow such as shear banding: In response to a homogeneous shear stress, part of the material becomes liquid and flows more and more easily with time, while the rest remains solid. Shear banding is an important phenomenon to understand and control when handling YSFs industrially. Now, Sébastien Manneville, of the École Normale Supérieure de Lyon, and colleagues have observed unexpected shear banding in a non-thixotropic, or “simple,” YSF. Unlike the thixotropic YSFs, which show shear banding in the steady state, the simple YSF's shear banding was transient—but the transient regime lasted a surprisingly long time.²

Stressed out

Complex fluids owe their complexity to structural elements, such as colloidal

particles, that are much larger than small molecules but much smaller than the bulk. Thixotropy can arise in a complex fluid (which could be a YSF or not) when those structures attract each other

to form aggregates that stiffen the material. When the fluid is forced to flow, the aggregates break down. As a result, more stress is required to initiate flow in a thixotropic YSF than to sustain it,




Figure 1. (a) Geometry of the Couette cell used in the experiment. Carbopol gel fills the gap between two concentric cylinders, and the inner cylinder rotates at a constant rate. The inset shows a velocity profile corresponding to homogeneous flow. (Adapted from ref. 2.) **(b)** Three velocity profiles recorded for a shear rate of 0.7 s^{-1} (a rotational period of about three minutes). After 27 s, flow is mostly confined to a thin shear band around the inner cylinder; after 1730 s, the shear band is thicker; and after 2250 s, the flow is nearly homogeneous. (Adapted from ref. 2.)

and the fluids don't always flow homogeneously, even when subjected to homogeneous stress.

But simple YSFs have no reason not to flow homogeneously. So Manneville and colleagues didn't expect to see any shear banding. When they set out to study their material's flow under shear stress, they were hoping to get some insight into how a YSF yields, or turns from solid to liquid. More specifically, they wanted to see whether the smoothness or roughness of the container walls had any effect.

The material they studied was carbopol gel, a main ingredient in hair gel and many pharmaceutical gels. Synthesized from carbopol powder and water, the gel itself is an assembly of soft, swollen polymer blobs. The researchers used a rheometer equipped with a Couette cell, shown in figure 1a: A layer of gel filled the 1.1-mm gap between two concentric cylinders, and the inner cylinder was rotated at a constant rate.

To measure the gel's velocity profile, the researchers used ultrasonic speckle velocimetry, a technique Manneville had developed for cheaply studying optically opaque fluids.³ They seeded the gel with micron-sized glass spheres, which scatter acoustic waves, and aimed an ultrasonic transducer at the sample at an angle to the direction of flow. Every millisecond or so, they shot an ultrasound pulse through the gel and recorded its reflection. From correlations between consecutive shots, they deduced the gel's velocity as a function of position and time.

Not so simple

Figure 1b shows some results from a typical run, during which the inner cylinder rotated once every three minutes. At first the gel flowed only near the moving wall, but with time the




Figure 2. Power-law dependence of the fluidization time (the duration of the transient shear banding) on the shear rate. Symbols represent different gap widths and different boundary conditions, as indicated by the legend. The gray line is a power-law fit with slope -2.3 . (Adapted from ref. 2.)

flowing band gradually widened and then suddenly gave way to homogeneous flow. That so-called fluidization time was surprisingly long: more than half an hour for the experiment shown.

Manneville and colleagues repeated the experiment, rotating the cylinder at different speeds. Not surprisingly, fluidization happened faster when the cylinder was spun faster. But quantitatively, the dependence took an unexpected form: The fluidization time was inversely proportional to the cylinder's speed raised to the 2.3 power. When the researchers looked at thicker layers of gel, they found a power-law dependence on the shear rate (the inner wall's linear speed divided by the gap width), as shown in figure 2. And it mattered little whether the inner and outer walls were bare plexiglass or whether they were coated with sandpaper to make them rough.

Where the exponent 2.3 comes from is unclear. The researchers think it's a material property, since using carbopol gel from a different batch—which may have had different-sized polymer blobs—was enough to change the exponent from 2.3 to about 3. They're now looking at gels prepared with different concentrations of carbopol powder to better understand

the material dependence. Extending the study to different simple YSFs, such as foams, would be more difficult: The high acoustic contrast between a foam's gas and liquid phases makes the ultrasound probe unsuitable.

The observation of transient shear banding shows that nonthixotropic YSFs are not as simple as they've been assumed to be; they don't yield uniformly, even under uniform stress. (The researchers briefly looked at other flow geometries to make sure that the shear banding was not an artifact of the slight stress heterogeneity induced by the circular geometry.) That the power-law relationship is independent of the layer thickness means that the fluid boundary doesn't simply diffuse across the gel. And the long fluidization times point to the importance of distinguishing between steady-state and transient regimes in future YSF models and experiments.

Johanna Miller

References

1. D. Bonn, M. M. Denn, *Science* **324**, 1401 (2009); P. C. F. Møller, J. Mewis, D. Bonn, *Soft Matter* **2**, 274 (2006).
2. T. Divoux et al., *Phys. Rev. Lett.* **104**, 208301 (2010).
3. S. Manneville, L. Bécu, A. Colin, *Eur. Phys. J.: Appl. Phys.* **28**, 361 (2004).

physics update

These items, with supplementary material, first appeared at <http://www.physicstoday.org>.

A thermometer for modern and extinct vertebrates. Robert Eagle of Caltech and his collaborators have shown that they can determine the body temperature of living and long-dead vertebrates by measuring the abundance of a molecule made of isotopes—an isotopologue—in bones, scales, and teeth. The isotopologue is a heavy version of the carbonate ion CO_3^{2-} . In a typical piece of bone or other biomineral, all but 1.8% of the CO_3^{2-} ions are made of the lightest carbon and oxygen isotopes, ^{12}C and ^{16}O . At around 45 ppm, $^{13}\text{C}^{18}\text{O}^{16}\text{O}_2^{2-}$ is barely present, but its scarcity is made up for by a useful property: The isotopologue's precise abundance depends on the ambient temperature when the biomineral first crystallized. The temperature dependence arises because lower temperatures boost the propensity of a heavy iso-

tope to form a bond with another heavy isotope rather than with a light isotope. Five years ago, Prosenjit Ghosh, who is now at the Indian Institute of Science, and his colleagues extracted CO_2 gas from carbonate crystals they'd made in the lab. From their measurements they derived a robust formula relating the abundance of $^{13}\text{C}^{18}\text{O}$ carbonate to its formation temperature. By applying the formula to tooth samples, Eagle could accurately predict the body temperature of five vertebrates, including the white rhino (37 °C) and the sand tiger shark (23 °C). From fossilized samples he could also predict the body temperature of the woolly mammoth (38 °C). Applying the paleothermometer to samples of other extinct vertebrates could reveal when vertebrates first became warm-blooded. (R. A. Eagle et al., *Proc. Natl. Acad. Sci. USA* **107**, 10377, 2010.)

—CD

An optical speed trap for Brownian motion. One hallmark of Albert Einstein's genius is his 1905 theory that the kinetic energy of pollen grains, dust, and other similarly sized objects in thermal